

Vanishing theorems of the basic harmonic forms on a complete foliated Riemannian manifold

Seoung Dal Jung and Huili Liu

Abstract. It is well-known that on a compact foliated Riemannian manifold $(M, \mathcal{F})$ with some transversal curvature conditions, there are no nontrivial basic harmonic r ($0 < r < q = \text{codim} \mathcal{F}$)-forms (M. Min-Oo et al., J. Reine Angew. Math. 415 (1991) [9]). In this paper, we extend the above facts to a complete foliated Riemannian manifold.

1 Introduction

Let $(M, g, \mathcal{F})$ be a foliated Riemannian manifold with a foliation $\mathcal{F}$ and a bundle-like metric g with respect to $\mathcal{F}$. A foliated Riemannian manifold is a Riemannian manifold with a Riemannian foliation, i.e., a foliation on a smooth manifold such that the normal bundle is endowed with a metric whose Lie derivative is zero along leaf directions (see [13]). A Riemannian metric on M is bundle-like if the leaves of the foliation $\mathcal{F}$ are locally equidistant, that is, the metric g on M induces a holonomy invariant transverse metric on the normal bundle $Q = TM/T\mathcal{F}$, where $T\mathcal{F}$ is the tangent bundle of $\mathcal{F}$. Every Riemannian foliation admits bundle-like metrics. Many researchers have studied basic forms and the basic Laplacian on foliated Riemannian manifolds. Basic forms are locally forms on the space of leaves; that is, forms ϕ satisfying $i(X)\phi = i(X)d\phi = 0$ for all $X \in T\mathcal{F}$. Basic forms are preserved by the exterior derivative and are used to define basic de-Rham cohomology groups $H_B^*(\mathcal{F})$. The basic Laplacian Δ_B for a given bundle-like metric is a version of the Laplace operator that preserves the basic forms. It is well-known [6,14] that on a closed oriented manifold M with a transversally oriented Riemannian foliation $\mathcal{F}$, $H_B^r(\mathcal{F}) \cong \mathcal{H}_B^r(\mathcal{F})$, where $\mathcal{H}_B^r(\mathcal{F}) = \ker \Delta_B$ is finite dimensional. And so $\chi_B(\mathcal{F}) = \sum_{r=0}^q (-1)^r \dim \mathcal{H}_B^r(\mathcal{F})$, where $\chi_B(\mathcal{F})$ is the basic Euler characteristic [2]. In 1991, M. Min-Oo et al. [9] proved that on a closed foliated Riemannian manifold M , if the transversal curvature operator of

2000 *Mathematics Subject Classification.* 53C12, 58E20

Key words and phrases. Basic harmonic forms, Basic Laplacian

$\mathcal{F}$ is positive definite, then $H_B^r(\mathcal{F}) = 0$ ($0 < r < q$), that is, any basic harmonic r -form is trivial.

In this paper, we study the basic r -forms on a complete foliated Riemannian manifold.

Main Theorem. *Let $(M, g, \mathcal{F})$ be a complete foliated Riemannian manifold and all leaves be compact. Assume that the mean curvature form is bounded and coclosed.*

(1) *If the transversal Ricci curvature of $\mathcal{F}$ is positive-definite, then any L^2 -basic harmonic 1-forms ϕ with $\phi \in \mathcal{S}_B$ are trivial.*

(2) *If the curvature endomorphism of $\mathcal{F}$ is positive-definite, then any L^2 -basic harmonic r -forms ϕ with $\phi \in \mathcal{S}_B$ are trivial.*

Here $\mathcal{S}_B$ is the Sobolev space of basic forms whose derivative belong to $L^2\Omega_B^*(\mathcal{F})$.

Note that in 1980, H. Kitahara [8] proved that under the same condition of the transversal Ricci curvature, there are no nontrivial basic Δ_T -harmonic 1-forms with finite global norms. Here Δ_T is a different operator to the basic Laplacian Δ_B . If $\mathcal{F}$ is minimal, then $\Delta_T = \Delta_B$.

2 Preliminaries

Let $(M, g, \mathcal{F})$ be a $(p + q)$ -dimensional complete foliated Riemannian manifold with a foliation $\mathcal{F}$ of codimension q and a bundle-like metric g with respect to $\mathcal{F}$. Let TM be the tangent bundle of M , $T\mathcal{F}$ its integrable subbundle given by $\mathcal{F}$, and $Q = TM/T\mathcal{F}$ the corresponding normal bundle of $\mathcal{F}$. Then we have an exact sequence of vector bundles

$$0 \longrightarrow T\mathcal{F} \longrightarrow TM \xrightarrow[\sigma]{\pi} Q \longrightarrow 0, \quad (2.1)$$

where $\pi : TM \rightarrow Q$ is a projection and $\sigma : Q \rightarrow T\mathcal{F}^\perp$ is a bundle map satisfying $\pi \circ \sigma = id$. Let g_Q be the holonomy invariant metric on Q induced by g , i.e., $\theta(X)g_Q = 0$ for any vector field $X \in T\mathcal{F}$, where $\theta(X)$ is the transverse Lie derivative [5]. Let R^Q and Ric^Q be the transversal curvature tensor and transversal Ricci operator of $\mathcal{F}$ with respect to the transversal Levi-Civita connection $\nabla^Q \equiv \nabla$ in Q [14], respectively. A differential form $\phi \in \Omega^r(M)$ is *basic* if $i(X)\phi = 0$ and $i(X)d\phi = 0$ for all $X \in T\mathcal{F}$. In a distinguished chart $(x_1, \dots, x_p; y_1, \dots, y_q)$ of

$\mathcal{F}$, a basic r -form ϕ is expressed by

$$\phi = \sum_{a_1 < \dots < a_r} \phi_{a_1 \dots a_r} dy_{a_1} \wedge \dots \wedge dy_{a_r},$$

where the functions $\phi_{a_1 \dots a_r}$ are independent of x . Let $\Omega_B^r(\mathcal{F})$ be the set of all basic r -forms on M . Then $\Omega^r(M) = \Omega_B^r(\mathcal{F}) \oplus \Omega_B^r(\mathcal{F})^\perp$ [1]. Now, we recall the star operator $\bar{*} : \Omega_B^r(\mathcal{F}) \rightarrow \Omega_B^{q-r}(\mathcal{F})$ given by [6,12]

$$\bar{*}\phi = (-1)^{p(q-r)} * (\phi \wedge \chi_{\mathcal{F}}), \quad \forall \phi \in \Omega_B^r(\mathcal{F}), \quad (2.2)$$

where $\chi_{\mathcal{F}}$ is the characteristic form of $\mathcal{F}$ and $*$ is the Hodge star operator associated to g . For any basic forms $\phi, \psi \in \Omega_B^r(\mathcal{F})$, it is well-known [12] that $\phi \wedge \bar{*}\psi = \psi \wedge \bar{*}\phi$ and $\bar{*}^2\phi = (-1)^{r(q-r)}\phi$. The operator d_B is the restriction of d to the basic forms, i.e., $d_B = d|_{\Omega_B^*(\mathcal{F})}$. Let $d_T = d_B - \kappa_B \wedge$ and $\delta_T = (-1)^{q(r+1)+1} \bar{*} d_B \bar{*}$, where κ_B is the basic part of the mean curvature form κ of $\mathcal{F}$ [1]. Note that κ_B is closed, i.e., $d\kappa_B = 0$ [10,14]. The operator $\delta_B : \Omega_B^r(\mathcal{F}) \rightarrow \Omega_B^{r-1}(\mathcal{F})$ is defined by

$$\delta_B \phi = (-1)^{q(r+1)+1} \bar{*} d_T \bar{*} \phi = \delta_T \phi + i(\kappa_B^\sharp) \phi, \quad (2.3)$$

where $(\cdot)^\sharp$ is the g_Q -dual vector field of $(\cdot)$. Generally, δ_B is not a restriction of δ on $\Omega_B^r(\mathcal{F})$, i.e., $\delta_B \neq \delta|_{\Omega_B^r(\mathcal{F})}$, where δ is the formal adjoint of d . But $\delta_B \omega = \delta \phi$ for any basic 1-form ϕ . Let $\Delta_B = d_B \delta_B + \delta_B d_B$ be a basic Laplacian. Then $\Delta^M|_{\Omega_B^0(\mathcal{F})} = \Delta_B$ [6], where Δ^M is the Laplacian on M . Let $\{E_a\} (a = 1, \dots, q)$ be a local orthonormal basic frame of Q and θ^a a g_Q -dual 1-form to E_a . We define $\nabla_{\text{tr}}^* \nabla_{\text{tr}} : \Omega_B^r(\mathcal{F}) \rightarrow \Omega_B^r(\mathcal{F})$ by

$$\nabla_{\text{tr}}^* \nabla_{\text{tr}} = - \sum_a \nabla_{E_a, E_a}^2 + \nabla_{\kappa_B^\sharp}, \quad (2.4)$$

where $\nabla_{X,Y}^2 = \nabla_X \nabla_Y - \nabla_{\nabla_X^M Y}$ for any $X, Y \in TM$ and ∇^M is the Levi-Civita connection with respect to g . Then the generalized Weitzenböck type formula on $\Omega_B^r(\mathcal{F})$ is given by [4]

$$\Delta_B \phi = \nabla_{\text{tr}}^* \nabla_{\text{tr}} \phi + F(\phi) + A_{\kappa_B^\sharp} \phi \quad (2.5)$$

for any $\phi \in \Omega_B^r(\mathcal{F})$, where $F = \sum_{a,b=1}^q \theta^a \wedge i(E_b) R^Q(E_b, E_a)$ and

$$A_Y \phi = \theta(Y) \phi - \nabla_Y \phi. \quad (2.6)$$

In particular, for a 1-form ϕ , $F(\phi)^\sharp = \text{Ric}^Q(\phi^\sharp)$ and $A_Y s = -\nabla_{\sigma(s)}\pi(Y)$. Let $\Omega_{B,o}^*(\mathcal{F})$ be the space of basic forms with compact supports.

Let ν be the transversal volume form, i.e., $*\nu = \chi_{\mathcal{F}}$. The pointwise inner product $\langle \cdot, \cdot \rangle$ on $\Omega_B^r(\mathcal{F})$ is given by

$$\langle \phi, \psi \rangle \nu = \phi \wedge \bar{*}\psi \quad (2.7)$$

for any basic forms $\phi, \psi \in \Omega_B^r(\mathcal{F})$. And the global inner product $\ll \cdot, \cdot \gg$ on $\Omega_B^*(\mathcal{F})$ is defined by

$$\ll \phi, \psi \gg = \int_M \langle \phi, \psi \rangle \mu_M \quad (2.8)$$

for any $\phi, \psi \in \Omega_B^r(\mathcal{F})$, one of which has compact support, where $\mu_M = \nu \wedge \chi_{\mathcal{F}}$ is the volume form with respect to g . It is well-known [4] that $\ll \nabla_{\text{tr}}^* \nabla_{\text{tr}} \phi, \psi \gg = \ll \nabla_{\text{tr}} \phi, \nabla_{\text{tr}} \psi \gg$ for any $\phi, \psi \in \Omega_{B,0}^r(\mathcal{F})$ and

$$\ll d_B \phi, \psi \gg = \ll \phi, \delta_B \psi \gg \quad (2.9)$$

for any $\phi \in \Omega_{B,o}^r(\mathcal{F})$, $\psi \in \Omega_{B,o}^{r+1}(\mathcal{F})$. The basic form ϕ is said to be L^2 -basic form if ϕ has finite global norm, i.e., $\|\phi\|^2 < \infty$. Let $\mathcal{H}_{B,2}^r(\mathcal{F})$ be the space of L^2 -basic harmonic forms, i.e.,

$$\mathcal{H}_{B,2}^r(\mathcal{F}) = \{\phi \in L^2 \Omega_B^r(\mathcal{F}) \mid d_B \phi = \delta_B \phi = 0\}. \quad (2.10)$$

Generally, the space $\mathcal{H}_{B,2}^r(\mathcal{F})$ can have infinite dimension. And if the dimension of $\mathcal{H}_{B,2}^r(\mathcal{F})$ is finite, then it depends on the bundle-like metric. Trivially, if M is compact, then $\mathcal{H}_{B,2}^r(\mathcal{F}) \cong H_B^r(\mathcal{F})$. So we study the vanishing properties of the L^2 -basic harmonic spaces on a complete foliated Riemannian manifold.

Remark 2.1 (1) The operator δ_T is the formal adjoint of d_B with respect to the global norm $(\cdot, \cdot)$, which is given by

$$(\phi, \psi) = \int_M \langle \phi, \psi \rangle \nu \wedge dx_1 \wedge \cdots \wedge dx_p. \quad (2.11)$$

Let $\Delta_T = d_B \delta_T + \delta_T d_B$ be a Laplacian. If the foliation is minimal, then $\delta_T = \delta_B$. So $\Delta_B = \Delta_T$.

(2) In 1980, H. Kitahara [8] proved that if the transversal Ricci curvature is nonnegative and positive at some point, then there are no nontrivial L^2 -basic Δ_T -harmonic 1-forms.

3 Vanishing theorem

Let $(M, g, \mathcal{F})$ be a complete foliated Riemannian manifold with a foliation $\mathcal{F}$ of codimension q and a bundle-like metric g with respect to $\mathcal{F}$. Assume that all leaves of $\mathcal{F}$ are compact. Now, we consider a smooth function μ on $\mathbb{R}$ satisfying

$$(i) \ 0 \leq \mu(t) \leq 1 \text{ on } \mathbb{R}, \quad (ii) \ \mu(t) = 1 \text{ for } t \leq 1, \quad (iii) \ \mu(t) = 0 \text{ for } t \geq 2.$$

Let x_0 be a point in M . For each point $y \in M$, we denote by $\rho(y)$ the distance between leaves through x_0 and y . For any real number $l > 0$, we define a Lipschitz continuous function ω_l on M by

$$\omega_l(y) = \mu(\rho(y)/l).$$

Trivially, ω_l is a basic function. Let $B(l) = \{y \in M | \rho(y) \leq l\}$ for $l > 0$. Then ω_l satisfies the following properties:

$$\begin{aligned} 0 \leq \omega_l(y) \leq 1 & \quad \text{for any } y \in M \\ \text{supp } \omega_l \subset B(2l) & \\ \omega_l(y) = 1 & \quad \text{for any } y \in B(l) \\ \lim_{l \rightarrow \infty} \omega_l = 1 & \\ |d_B \omega_l| \leq \frac{C}{l} & \quad \text{almost everywhere on } M, \end{aligned}$$

where C is a positive constant independent of l [15]. Hence $\omega_l \psi$ has compact support for any basic form $\psi \in \Omega_B^*(\mathcal{F})$ and $\omega_l \psi \rightarrow \psi$ (strongly) when $l \rightarrow \infty$.

Lemma 3.1 [7] *For any $\phi \in \Omega_B^r(\mathcal{F})$, there exists a number A depending only on μ , such that*

$$\begin{aligned} \|d_B \omega_l \wedge \phi\|_{B(2l)}^2 &\leq \frac{qA^2}{l^2} \|\phi\|_{B(2l)}^2, \\ \|d_B \omega_l \wedge \bar{*}\phi\|_{B(2l)}^2 &\leq \frac{qA^2}{l^2} \|\phi\|_{B(2l)}^2, \\ \|d_B \omega_l \otimes \phi\|_{B(2l)}^2 &\leq \frac{qA^2}{l^2} \|\phi\|_{B(2l)}^2, \end{aligned}$$

where $\|\phi\|_{B(2l)}^2 = \ll \phi, \phi \gg_{B(2l)} = \int_{B(2l)} \langle \phi, \phi \rangle \mu_M$.

Proposition 3.2 *For any L^2 -basic form ψ , if $\Delta_B \psi = 0$, then $d_B \psi = 0$ and $\delta_B \psi = 0$.*

Proof. Let ψ be a L^2 -basic form. Then we have

$$\ll \Delta_B \psi, \omega_l^2 \psi \gg_{B(2l)} = \ll d_B \psi, d_B(\omega_l^2 \psi) \gg_{B(2l)} + \ll \delta_B \psi, \delta_B(\omega_l^2 \psi) \gg_{B(2l)}. \quad (3.1)$$

By a direct calculation, we have

$$d_B(\omega_l^2 \psi) = \omega_l^2 d_B \psi + 2\omega_l d_B \omega_l \wedge \psi, \quad (3.2)$$

$$\delta_B(\omega_l^2 \psi) = \omega_l^2 \delta_B \psi + (-1)^{q(r+1)+1} \bar{*}(2\omega_l d_B \omega_l \wedge \bar{*}\psi). \quad (3.3)$$

From (3.1), (3.2) and (3.3), if $\Delta_B \psi = 0$, then

$$\begin{aligned} & \|\omega_l d_B \psi\|_{B(2l)}^2 + \|\omega_l \delta_B \psi\|_{B(2l)}^2 \\ &= -2 \ll \omega_l d_B \psi, d_B \omega_l \wedge \psi \gg_{B(2l)} + 2(-1)^{q(r+1)} \ll \omega_l \delta_B \psi, \bar{*}(d_B \omega_l \wedge \bar{*}\psi) \gg_{B(2l)}. \end{aligned}$$

Hence by the Schwartz's inequality and Lemma 3.1, we have

$$\begin{aligned} & \|\omega_l d_B \psi\|_{B(2l)}^2 + \|\omega_l \delta_B \psi\|_{B(2l)}^2 \\ & \leq \epsilon_1 \|\omega_l d_B \psi\|_{B(2l)}^2 + \epsilon_2 \|\omega_l \delta_B \psi\|_{B(2l)}^2 + \frac{B_1}{l} \|\psi\|_{B(2l)}^2 \end{aligned}$$

for some positive real numbers ϵ_1, ϵ_2 and B_1 . Therefore, we have

$$\|\omega_l d_B \psi\|_{B(2l)}^2 + \|\omega_l \delta_B \psi\|_{B(2l)}^2 \leq \frac{B_2}{l} \|\psi\|_{B(2l)}^2$$

for some positive real number B_2 . Since ψ is the L^2 -basic form, letting $l \rightarrow \infty$, $d_B \psi = \delta_B \psi = 0$. $\square$

Remark 3.3 In 1979, H. Kitahara [7] proved the corresponding result with the Laplacian Δ_T . Namely, on a complete foliated manifold, if $\Delta_T \phi = 0$, then $d_B \phi = \delta_T \phi = 0$.

Now we prove the vanishing theorem of the L^2 -basic harmonic form on a complete foliated Riemannian manifold. First of all, we prepare some lemmas.

Lemma 3.4 *Let $(M, g, \mathcal{F})$ be a complete foliated Riemannian manifold whose leaves are compact. Suppose that κ_B is bounded and coclosed. Then for any L^2 -basic harmonic form ϕ ,*

$$\limsup_{l \rightarrow \infty} \ll A_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)} = 0. \quad (3.4)$$

Proof. Let ϕ be a L^2 -basic harmonic form. Since $\theta(X)\phi = d_B i(X)\phi$, from (2.6) we have

$$\ll A_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)} = \ll d_B i(\kappa_B^\#) \phi, \omega_l^2 \phi \gg_{B(2l)} - \ll \nabla_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)} \quad (3.5)$$

Since $\delta_B \phi = 0$, from (3.3) and Lemma 3.1, we have

$$\begin{aligned} |\ll d_B i(\kappa_B^\#) \phi, \omega_l^2 \phi \gg_{B(2l)}| &= 2 |\ll \omega_l i(\kappa_B^\#) \phi, \bar{*}(d_B \omega_l \wedge \bar{*} \phi) \gg_{B(2l)}| \\ &\leq \epsilon_3 \|\omega_l i(\kappa_B^\#) \phi\|_{B(2l)}^2 + \frac{1}{\epsilon_3} \|d_B \omega_l \wedge \bar{*} \phi\|_{B(2l)}^2 \\ &\leq \epsilon_3 \|\omega_l i(\kappa_B^\#) \phi\|_{B(2l)}^2 + \frac{B_3}{l^2} \|\phi\|_{B(2l)}^2 \end{aligned}$$

for some positive real numbers ϵ_3 and B_3 . By using $|i(\kappa_B^\#) \phi|^2 + |\kappa_B \wedge \phi|^2 = |\kappa_B|^2 |\phi|^2$, we have

$$|\ll d_B i(\kappa_B^\#) \phi, \omega_l^2 \phi \gg_{B(2l)}| \leq \epsilon_3 \max(|\kappa_B|^2) \|\omega_l \phi\|_{B(2l)}^2 + \frac{B_3}{l^2} \|\phi\|_{B(2l)}^2 \quad (3.6)$$

On the other hand, since $\delta_B \kappa_B = 0$, by a direct calculation, we have

$$\begin{aligned} \ll \nabla_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)} &= \frac{1}{2} \ll d_B (|\omega_l \phi|^2), \kappa_B \gg_{B(2l)} - \ll \omega_l \phi, \kappa_B^\# (\omega_l) \phi \gg_{B(2l)} \\ &= - \ll \omega_l \phi, \kappa_B^\# (\omega_l) \phi \gg_{B(2l)}. \end{aligned}$$

Hence by the Schwartz inequality, we have

$$\begin{aligned} |\ll \nabla_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)}| &= |\ll \omega_l \phi, \kappa_B^\# (\omega_l) \phi \gg_{B(2l)}| \\ &\leq \epsilon_4 \|\omega_l \phi\|_{B(2l)}^2 + \frac{B_4}{l^2} \max(|\kappa_B|^2) \|\phi\|_{B(2l)}^2 \end{aligned} \quad (3.7)$$

for a positive real numbers ϵ_4 and B_4 . From (3.6) and (3.7), by letting $l \rightarrow \infty$, we have

$$\begin{aligned} \limsup_{l \rightarrow \infty} |\ll d_B i(\kappa_B^\#) \phi, \omega_l^2 \phi \gg_{B(2l)}| &\leq \epsilon_3 \max(|\kappa_B|^2) \|\phi\|^2, \\ \limsup_{l \rightarrow \infty} |\ll \nabla_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)}| &\leq \epsilon_4 \|\phi\|^2. \end{aligned}$$

Since ϵ_3 and ϵ_4 are arbitrary positive numbers, we have

$$\limsup_{l \rightarrow \infty} |\ll d_B i(\kappa_B^\#) \phi, \omega_l^2 \phi \gg_{B(2l)}| = 0, \quad (3.8)$$

$$\limsup_{l \rightarrow \infty} |\ll \nabla_{\kappa_B^\#} \phi, \omega_l^2 \phi \gg_{B(2l)}| = 0. \quad (3.9)$$

Hence from (3.5), (3.8) and (3.9), the proof is completed. $\square$

Theorem 3.5 *Let $(M, g, \mathcal{F})$ be as in Lemma 3.4. Suppose that κ_B is bounded and coclosed. If the curvature endomorphism F of $\mathcal{F}$ is positive-definite, then any L^2 -basic harmonic r -forms ϕ with $\phi \in \mathcal{S}_B$ are trivial, i.e., $\mathcal{H}_{B,2}^r(\mathcal{F}) = \{0\}$.*

Proof. Let ϕ be a L^2 -basic harmonic r -form. From (2.5) and Proposition 3.2, we have

$$\langle \nabla_{\text{tr}}^* \nabla_{\text{tr}} \phi, \omega_l^2 \phi \rangle + \langle F(\phi), \omega_l^2 \phi \rangle + \langle A_{\kappa_B^\sharp} \phi, \omega_l^2 \phi \rangle = 0. \quad (3.10)$$

On the other hand, a direct calculation gives

$$\ll \nabla_{\text{tr}}^* \nabla_{\text{tr}} \phi, \omega_l^2 \phi \gg_{B(2l)} = \ll \nabla_{\text{tr}} \phi, 2\omega_l d_B \omega_l \otimes \phi \gg_{B(2l)} + \|\omega_l \nabla_{\text{tr}} \phi\|_{B(2l)}^2. \quad (3.11)$$

From Lemma 3.1, we have

$$| \ll \nabla_{\text{tr}} \phi, 2\omega_l d_B \omega_l \otimes \phi \gg_{B(2l)} | \leq \epsilon_5 \|\omega_l \nabla_{\text{tr}} \phi\|_{B(2l)}^2 + \frac{B_5}{l^2} \|\phi\|_{B(2l)}^2$$

for some positive constants ϵ_5 and B_5 . Hence by letting $l \rightarrow \infty$, we have

$$\limsup_{l \rightarrow \infty} \ll \nabla_{\text{tr}} \phi, 2\omega_l d_B \omega_l \otimes \phi \gg_{B(2l)} \leq \epsilon_5 \|\nabla_{\text{tr}} \phi\|^2.$$

Since ϵ_5 is arbitrary and $\phi \in \mathcal{S}_B$ (i.e., $\|\nabla_{\text{tr}} \phi\|^2 < \infty$), we have

$$\limsup_{l \rightarrow \infty} \ll \nabla_{\text{tr}} \phi, 2\omega_l d_B \omega_l \otimes \phi \gg_{B(2l)} = 0. \quad (3.12)$$

Hence from (3.11), (3.12) and Lemma 3.4, we have

$$\|\nabla_{\text{tr}} \phi\|^2 + \limsup_{l \rightarrow \infty} \ll F(\phi), \omega_l^2 \phi \gg_{B(2l)} = 0, \quad (3.13)$$

which complete the proof. $\square$

Since $F(\phi^\sharp) = \text{Ric}^Q(\phi^\sharp)$ for any basic 1-form ϕ , we have the following corollary.

Corollary 3.6 *Let $(M, g, \mathcal{F})$ be as in Lemma 3.4. Suppose that κ_B is bounded and coclosed. If the transversal Ricci curvature Ric^Q is positive-definite, then any L^2 -basic harmonic 1-forms ϕ with $\phi \in \mathcal{S}_B$ are trivial, $\mathcal{H}_{B,2}^1(\mathcal{F}) = \{0\}$.*

Acknowledgements. The first author was supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government (MSIP) (NRF-2015R1A2A2A01003491) and the second author was supported by NSFC (No. 11371080).

References

- [1] J. A. Alvarez López, *The basic component of the mean curvature of Riemannian foliations*, Ann. Global Anal. Geom. 10 (1992), 179-194.
- [2] V. Blefi, E. Park and K. Richardson, *A Hopf index theorem for foliations*, Diff. Geom. Appl. 18 (2003), 319-341.
- [3] J. J. Hebda, *Curvature and focal points in Riemannian foliations*, Indiana Univ. Math. J. 35 (1986), 321-331.
- [4] S. D. Jung, *The first eigenvalue of the transversal Dirac operator*, J. Geom. Phys. 39 (2001), 253-264.
- [5] F. W. Kamber and Ph. Tondeur, *Infinitesimal automorphisms and second variation of the energy for harmonic foliations*, Tôhoku Math. J. 34 (1982), 525-538.
- [6] F. W. Kamber and Ph. Tondeur, *De Rham-Hodge theory for Riemannian foliations*, Math. Ann. 277 (1987), 415-431.
- [7] H. Kitahara, *Remarks on square-integrable basic cohomology spaces on a foliated Riemannian manifold*, Kodai Math. J. 2 (1979), 187-193.
- [8] H. Kitahara, *Nonexistence of nontrivial $\square''$ -harmonic 1-forms on a complete foliated Riemannian manifold*, Trans. Amer. Math. Soc. 262 (1980), 429-435.
- [9] M. Min-Oo, E. Ruh and Ph. Tondeur, *Vanishing theorems for the basic cohomology of Riemannian foliations*, J. Reine Angew. Math. 415 (1991), 167-174.
- [10] J. S. Pak and S. D. Jung, *A transversal Dirac operator and some vanishing theorems on a complete foliated Riemannian manifold*, Math. J. Toyama Univ. 16 (1993), 97-108.
- [11] J. S. Pak and S. D. Jung, *Some vanishing theorems on complete Kähler foliations*, Acta Math. Hungar. 77 (1997), 15-28.
- [12] E. Park and K. Richardson, *The basic Laplacian of a Riemannian foliation*, Amer. J. Math. 118 (1996), 1249-1275.
- [13] B. Reinhart, *Differential Geometry of Foliations*, Springer-Verlag, 1983.

- [14] Ph. Tondeur, *Geometry of foliations*, Basel: Birkhäuser Verlag, 1997.
- [15] S. Yorozu, *Notes on square-integrable cohomology spaces on certain foliated manifolds*, Trans. Amer. Math. Soc. 255 (1979), 329-341.

Department of Mathematics and Research Institute for Basic Sciences, Jeju National University, Jeju 690-756, Korea
E-mail address : sdjung@jejunu.ac.kr

Department of Mathematics, Northeastern University, Shenyang 110004, P. R. China
E-mail address : liuhl@mail.neu.edu.cn